

Quantum corrections to spin effects in general relativity

G.G. Kirilin¹

Budker Institute of Nuclear Physics
630090 Novosibirsk, Russia

Abstract

Quantum power corrections to the gravitational spin-orbit and spin-spin interactions, as well as to the Lense-Thirring effect, were found for particles of spin $1/2$. These corrections arise from diagrams of second order in Newton gravitational constant G with two massless particles in the unitary cut in the t -channel. The corrections obtained differ from the previous calculation of the corrections to spin effects for rotating compound bodies with spinless constituents.

PACS: 04.60.-m

¹g_kirilin@mail.ru

1 Introduction

It is common knowledge that the scattering amplitude $\mathcal{M} = \langle 1(p_1 - q), 2(p_2 + q) | 1(p_1), 2(p_2) \rangle$ of two gravitating bodies contains terms logarithmic in the momentum transfer $\log |q^2|$ in the $q^2 \rightarrow 0$ limit. The terms arise from diagrams with two massless particles in the unitary cut in the t-channel. In the nonrelativistic limit these terms generate quantum power corrections to the classical Hamiltonian [8]. Thus quantum corrections to the Newton potential take the form [1-20]

$$U = -\frac{41}{10\pi} \frac{G^2 \hbar m_1 m_2}{c^3 r^3} - \frac{4}{15\pi} \frac{G^2 \hbar m_1 m_2}{c^3 r^3} - \frac{N_\nu}{15\pi} \frac{G^2 \hbar m_1 m_2}{c^3 r^3}. \quad (1)$$

The first term in the expression (1) corresponds to the contribution of gravitons, the second term corresponds to the contribution of photons and the last one is the contribution of neutrinos, where N_ν is the number of massless two-component neutrinos (we put $c = 1$, $\hbar = 1$ below).

In the case of interaction of scalar particles, all contributions logarithmic in q^2 can be separated into two parts. The first one is the contributions in which the coefficient before $\log |q^2|$ is a function of $s = (p_1 + p_2)^2$ with cuts in the complex plane. These contributions can be generated by "box" diagrams only (see Figs. 5 c, d) that have unitary cut in the s - or u -channel. These contributions, called in [17] irreducible, can be calculated by the double dispersion relation in variables s and $t = q^2$. The distinctive characteristic of the contributions is that they are infrared divergent (see expression (70)) and, correspondingly, contain logarithms where q^2 plays a role of an ultraviolet cutoff $\log |q^2|/\lambda^2$. The second part is the contributions that are entire rational functions of s . They arise from the diagrams that have no unitary cut in the s - or u -channel and also from the reducible part of the "box" diagrams. These contributions contain logarithms with the transfer momentum as an infrared cutoff, for example, logarithms that contain ultraviolet cutoff $\log \Lambda^2/|q^2|$ or the masses of the interacting particles $\log m_i^2/|q^2|$.

As shown in [20], the second part of the amplitude can be represented as a matrix element of an effective nonlocal operator of interaction of two energy-momentum tensors (i.e., an effective graviton exchange) in the $q^2 \rightarrow 0$ limit (but for arbitrary s):

$$\hat{\mathcal{L}}_0 = \sqrt{-g} G^2 \log |q^2| \left(\frac{23}{5} T_{\mu\nu} T^{\mu\nu} - \frac{31}{30} T^2 \right) = \log |q^2| \frac{\sqrt{-g}}{(8\pi)^2} \left(\frac{23}{5} R_{\mu\nu} R^{\mu\nu} - \frac{31}{30} R^2 \right), \quad (2)$$

where $T_{\mu\nu}$, $R_{\mu\nu}$ are the energy-momentum tensor of the matter and the Ricci tensor, respectively. The main feature of the operator is that all descriptions of the interacting particles appear only in the form of a *product* of the energy-momentum tensors, whereas the "propagator" of the effective graviton is independent of the type of interacting particles². In contrast with the operator (2), the irreducible contributions of the diagrams 5 c, d contain nonmultiplicative dependence on momenta and masses of the particles and cannot be represented as operators like (2).

On the basis of this operator, power quantum corrections to the Schwarzschild metric were calculated in [20]. Corrections to the interactions dependent on internal angular momentum (spin) of a compound body were also considered. There are three ways to calculate these corrections: first of all, it is possible to calculate quantum corrections to the Kerr metric for rotating compound body and to use covariant spin equation of motion in an external gravitational field [21] as performed in [20]; the second way is to integrate velocity dependent interactions,

²Recall that here we are dealing with scalar particles only.

also calculated in [20], over the volumes of the interacting bodies; the last one is to put spin dependent energy-momentum tensor to the operator (2) explicitly.

The purpose of this work is to calculate power quantum corrections to the spin-dependent interactions for particles of spin 1/2. It was shown in an elementary way [20] that spin-averaged contributions containing q^2 as an infrared cutoff are exactly the same as (2). The irreducible parts of the diagrams 5 c, d averaged over spins coincide with the scalar ones since the logarithm of the infrared cutoff $\log \lambda$ is cancelled by radiation of long-wave gravitons which is spin-independent. However, it was noticed that the spin effects for rotating compound bodies with scalar constituents can differ from the ones for particles of spin 1/2. The reason is that, apart from the operator (2), the leading contribution to the spin effects can be generated by operators of higher order in q/m . For example, first order in q/m operators have the form:

$$\begin{aligned}\hat{O}_1 &= \sqrt{-g} \log |q^2| \frac{G}{16\pi} R^{\mu\nu}{}_{\alpha\beta;\mu} t_{(a)}^\alpha t_{(b)}^\beta \left[\sum_i \frac{1}{4m_i} \bar{\psi}_i \left(-i \overleftarrow{D}_\nu \Sigma^{ab} + \Sigma^{ab} i \overrightarrow{D}_\nu \right) \psi_i \right] \\ &= \sqrt{-g} \log |q^2| \frac{G}{16\pi} (R^\nu{}_{\beta;\alpha} - R^\nu{}_{\alpha;\beta}) t_{(a)}^\alpha t_{(b)}^\beta \left[\sum_i \frac{1}{4m_i} \bar{\psi}_i \left(-i \overleftarrow{D}_\nu \Sigma^{ab} + \Sigma^{ab} i \overrightarrow{D}_\nu \right) \psi_i \right],\end{aligned}\quad (3)$$

$$\hat{O}_2 = \sqrt{-g} \log |q^2| \frac{G^2}{2} (T^\nu{}_{\beta;\alpha} - T^\nu{}_{\alpha;\beta}) t_{(a)}^\alpha t_{(b)}^\beta \left[\sum_i \frac{1}{4m_i} \bar{\psi}_i \left(-i \overleftarrow{D}_\nu \Sigma^{ab} + \Sigma^{ab} i \overrightarrow{D}_\nu \right) \psi_i \right], \quad (4)$$

where left-hand-side and right-hand-side derivatives are defined as

$$\overrightarrow{D}_\alpha = \overrightarrow{\partial}_\alpha - \frac{i}{2} \Sigma^{ab} \gamma_{abc} t_\alpha^{(c)}, \quad \overleftarrow{D}_\alpha = \overleftarrow{\partial}_\alpha + \frac{i}{2} \Sigma^{ab} \gamma_{abc} t_\alpha^{(c)}. \quad (5)$$

Here $\Sigma^{ab} = \frac{i}{2} \sigma^{ab}$ are the Lorentz group generators, γ_{abc} are the Ricci rotation coefficients, $t_{(a)}^\mu$ is a tetrad of basis vectors that are specified by relation $t_{(a)}^\mu t_{\mu(b)} = \eta_{ab}$, where η_{ab} is the metric tensor of flat space-time. Taking into account Einstein and Dirac equations in an external gravitational field

$$R_{\mu\nu} = 8\pi G \left(T_{\mu\nu} - \frac{1}{2} g_{\mu\nu} T \right), \quad \left(i t_b^\nu \gamma^b \overrightarrow{D}_\nu - m \right) \psi = 0, \quad (6)$$

note that the difference of the operators $\hat{O}_1$ and $\hat{O}_2$ is an operator of second order in q/m

$$\hat{O}_1 - \hat{O}_2 = \sqrt{-g} \log |q^2| \frac{G^2}{16} T \left(\sum_i \frac{\square}{m_i^2} T_i \right) + \text{total derivative}$$

and makes zero contribution to the spin-dependent interactions.

Besides, contributions to the spin-spin interactions can be generated by operators of the second order in q/m :

$$\hat{O}_3 = \sqrt{-g} \log |q^2| D_\mu \left(t_{(a)}^\mu \bar{\psi}_2 \gamma^5 \gamma^a \psi_2 \right) D_\nu \left(t_{(b)}^\nu \bar{\psi}_1 \gamma^5 \gamma^b \psi_1 \right), \quad (7)$$

$$\hat{O}_4 = \sqrt{-g} \log |q^2| D^\mu \left(t_{(a)}^\nu \bar{\psi}_2 \gamma^5 \gamma^a \psi_2 \right) D_\mu \left(t_{\nu(b)} \bar{\psi}_1 \gamma^5 \gamma^b \psi_1 \right). \quad (8)$$

The plan of this paper is as follows: in Section 2 we describe useful Feynman rules with effective vertices introduced in [20] and also list possible types of spin-dependent amplitudes

with arbitrary coefficients (form factors). The explicit expressions for all the form factors in the $t \rightarrow 0$ limit are presented in Section 3. Results and conclusions are given in Sections 4 and 5.

2 Effective vertices




Figure 1. Vertices




Figure 2. Compton scattering amplitude

The single-graviton vertex for a spinor particle is (see Fig. 1 a)

$$V_{\mu\nu}^{(1/2)} = -\frac{i\kappa}{4} \bar{\psi}(p') \left\{ I_{\mu\nu, \alpha\beta} P^\alpha \gamma^\beta - \eta_{\mu\nu} (\hat{P} - 2m) \right\} \psi(p), \quad (9)$$

here $\kappa = \sqrt{32\pi G}$, $P = p + p'$, $I_{\mu\nu}^{\alpha\beta} = \frac{1}{2} (\delta_\mu^\alpha \delta_\nu^\beta + \delta_\mu^\beta \delta_\nu^\alpha)$ is a sort of a unit operator with the property

$$I_{\mu\nu}^{\alpha\beta} t_{\alpha\beta} = t_{\mu\nu}$$

for any symmetric tensor $t_{\alpha\beta}$. The last term in braces, expression (9), proportional to the Minkowski metric tensor $\eta_{\mu\nu}$ vanishes on-mass-shell. It is convenient to exclude such terms from the diagrams of Compton scattering amplitude containing a pole in the s - or u -channel (see Fig. 2). This term cancels fermion propagator and generates an additional contribution to the contact interaction of a mass-shell fermion with two gravitons (see Fig. 1 b). On rearrangement, the double graviton vertex has a form:

$$V_{\kappa\lambda, \rho\sigma}^{(1/2)} = i \frac{3}{4} \kappa^2 I_{\kappa\lambda, \alpha\delta} I_{\beta}^{\delta, \rho\sigma} T_{(1/2)}^{\alpha\beta}, \quad (10)$$

where $T_{\alpha\beta}^{(1/2)}$ is the energy-momentum tensor of the fermion field

$$T_{\mu\nu}^{(1/2)} = \frac{1}{2} I_{\mu\nu, \alpha\beta} P^\alpha \bar{u}(p') \gamma^\beta \psi(p). \quad (11)$$

For further Feynman rules the reader is referred to the papers [18], [20].

The rest of the pole diagrams (Fig.2) is proportional to the Compton scattering amplitude in quantum electrodynamics. For example, consider diagram shown in Fig. 2 a

$$M_{\alpha\beta, \gamma\delta} = \frac{i \kappa^2}{16(l^2 - m^2)} I_{\gamma\delta, \rho\sigma} I_{\alpha\beta, \phi\omega} (p' + l)^\rho (p + l)^\phi \left[\bar{u}(p') \gamma^\sigma (\hat{l} + m) \gamma^\omega u(p) \right]. \quad (12)$$

It is useful to decompose the amplitude of scattering of photon by an electron into independent spin structures:

$$M_{QED}^{\sigma\omega} = \bar{u}(p') \gamma^\sigma (\hat{l} + m) \gamma^\omega u(p) \\ = \frac{1}{2} [(P - K)^\sigma \delta_\rho^\omega + (P - K)^\omega \delta_\rho^\sigma - q^\sigma \delta_\rho^\omega + \delta_\rho^\sigma q^\omega + K_\rho \eta^{\sigma\omega}] J^\rho + \frac{i}{2} \epsilon^{\sigma\omega\lambda\eta} K_\lambda S_\eta, \quad (13)$$

here $P = 2p - q$, $K = 2k - q$, and J^μ , S^μ are the vector current and spin four-vector (axial current), respectively:

$$J^\mu = \bar{u}(p') \gamma^\mu u(p), \quad S^\mu = \bar{u}(p') \gamma_5 \gamma^\mu u(p), \quad \gamma_5 = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}. \quad (14)$$

Diagram shown in Fig. 2 b can be considered in a similar manner. Vector current J^μ can be also decomposed into a convective part and a part containing spin explicitly:

$$\left(1 - \frac{q^2}{4m^2}\right) J^\mu = \frac{P^\mu}{2m} \bar{u}(p') u(p) + \frac{i}{4m^2} \epsilon^{\mu\gamma\alpha\delta} q_\gamma P_\alpha S_\delta. \quad (15)$$

Therefore, all vertices and lines can be decomposed into two structure including bispinor amplitudes of a fermion, namely, $\bar{u}(p') u(p)$ and S^μ . Thus, matrix element of the fermion-fermion scattering can be separated into three groups. The first one does not include $S_\mu^{1,2}$ explicitly:

$$\mathcal{M}_N(q) = \mathcal{F}_N(s) G^2 m_1 m_2 (\bar{u}_1 u_1) (\bar{u}_2 u_2) \log |q^2|, \quad (16)$$

the second one contains the terms which are proportional to the first power of $S_\mu^{1,2}$:

$$\mathcal{M}_{SO}(q) = i \mathcal{F}_{SO}(s) G^2 \epsilon_{\alpha\beta\gamma\delta} q^\alpha P_1^\beta P_2^\gamma \left(\frac{S_2^\delta}{m_2} (\bar{u}_1 u_1) + \frac{S_1^\delta}{m_1} (\bar{u}_2 u_2) \right) \log |q^2|, \quad (17)$$

and the last part contains terms quadratic in spin:

$$\mathcal{M}_{SS} = G^2 \left(\mathcal{F}_{SS}^{(1)}(s) (q \cdot S_1) (q \cdot S_2) + \mathcal{F}_{SS}^{(2)}(s) t (S_1 \cdot S_2) \right) \log |q^2|. \quad (18)$$

In the first Born approximation corrections to a potential are given by a Fourier transform of nonrelativistic amplitude

$$U(\mathbf{r}) = \int \frac{d^3 \mathbf{q}}{(2\pi)^3} \mathcal{A}(\mathbf{q}) e^{-i\mathbf{q} \cdot \mathbf{r}} = - \int \frac{d^3 \mathbf{q}}{(2\pi)^3} e^{-i\mathbf{q} \cdot \mathbf{r}} \lim_{s \rightarrow (m_1 + m_2)^2} \lim_{t \rightarrow 0} \frac{\mathcal{M}}{\prod_i \sqrt{2E_i}}. \quad (19)$$

Since we are interested in amplitudes in the $t \rightarrow 0$ limit, the form factors $\mathcal{F}_i$ are the functions of the variable s . It is convenient to introduce a new variable

$$\tilde{\gamma} = (s - m_1^2 - m_2^2)/(2m_1 m_2) \quad (20)$$

which corresponds to the γ -factor of one of the interacting particles in the limit when the mass of the other goes to infinity.




Figure 3. Graviton-graviton interaction




Figure 4. Graviton-fermion interaction

3 Matrix elements

We start the discussion of loops with the graviton-graviton interaction shown in Fig. 3. It is easy to calculate this contribution using the effective operator derived in [7]

$$\mathcal{L}_{gg} = -\log |q^2| \frac{\sqrt{-g}}{(8\pi)^2} \left(\frac{21}{15} R^{\mu\nu} R_{\mu\nu} + \frac{1}{30} R^2 \right). \quad (21)$$

By substituting the Ricci tensor from the Einstein equations (6) one gets the following matrix elements:

$$\mathcal{F}_N(\tilde{\gamma}) = -\frac{1}{15} (42\tilde{\gamma}^2 + 1), \quad \mathcal{F}_{SO}(\tilde{\gamma}) = -\frac{7}{20} \tilde{\gamma}, \quad (22)$$

$$\mathcal{F}_{SS}^{(1)}(\tilde{\gamma}) = -\mathcal{F}_{SS}^{(2)}(\tilde{\gamma}) = \frac{7}{20} (1 - 2\tilde{\gamma}^2). \quad (23)$$

Diagrams containing the graviton-fermion interaction (see Fig. 4) make the following contribution to the matrix elements:

$$\mathcal{F}_N(\tilde{\gamma}) = 2 (6\tilde{\gamma}^2 + 1), \quad \mathcal{F}_{SO}(\tilde{\gamma}) = \frac{\tilde{\gamma}}{2}, \quad \mathcal{F}_{SS}^{(1)}(\tilde{\gamma}) = -\mathcal{F}_{SS}^{(2)}(\tilde{\gamma}) = \frac{1}{2} (1 - 2\tilde{\gamma}^2). \quad (24)$$

These corrections coincide with the corresponding form factor corrections calculated in [19]. By analogy with (21) the contribution of the diagrams including graviton-fermion interaction (Fig. 4) can be represented as an effective operator:

$$\mathcal{L}_{gf} = \sqrt{-g} \log |q^2| \frac{G}{4\pi} (3R^{\mu\nu} T_{\mu\nu} - 2RT) - 8\hat{O}_1, \quad (25)$$

where the operator $\hat{O}_1$ is defined by (3).

The rest of the diagrams (Fig. 5) corresponds to the fermion-fermion interaction. The sum of the diagrams Figs. 5 a, b makes the following contribution to the matrix elements:

$$\mathcal{F}_N(\tilde{\gamma}) = -3 (5\tilde{\gamma}^2 - 1), \quad \mathcal{F}_{SO}(\tilde{\gamma}) = -\frac{15}{16} \tilde{\gamma}, \quad \mathcal{F}_{SS}^{(1)}(\tilde{\gamma}) = -\mathcal{F}_{SS}^{(2)}(\tilde{\gamma}) = 0. \quad (26)$$

The sum of these diagrams does not contribute to the spin-spin interaction. The corresponding effective operator is




Figure 5. Fermion-fermion interaction

$$\mathcal{L}_{ff}^{(ab)} = -\sqrt{-g} \log |q^2| G^2 \left(\frac{15}{2} T^{\mu\nu} T_{\mu\nu} - \frac{3}{2} T^2 \right) + \frac{15}{2} \hat{O}_2, \quad (27)$$

where the operator $\hat{O}_2$ is defined by (4).

There are a number of points to be made before considering the contribution of the diagrams Figs. 5 c, d. As stated above, the analysis of these diagrams is complicated by the presence of infrared divergencies. Aside from the integrals (63)-(69) containing q^2 as an infrared cutoff, there are contributions generated by integrals (70), (72) containing q^2 as an ultraviolet cutoff. There is a unitary cut in the s - or u - channel in the last-mentioned integrals, and consequently, there is a nonpolynomial dependence on s . We present the results for the reducible and irreducible parts for each type of the matrix elements separately. However, as we shall see later, the reducible part also contains a nonpolynomial dependence on s . The reason is a singularity in the $\Delta \rightarrow 0$ limit in vector and tensor integrals of the $K_\mu^{(s)}$ type (see expression (76) in Appendix) where

$$\Delta = \frac{1}{4s} ((P_1 P_2)^2 - P_1^2 P_2^2), \quad P_1 = p_1 + p'_1, \quad P_2 = p_2 + p'_2. \quad (28)$$

In the center of mass frame the variable Δ takes the form:

$$\Delta = 4|\vec{p}|^2 \cos^2 \theta/2. \quad (29)$$

Terms proportional to $\log(\sin \theta/2)/(\cos^2 \theta/2)$ are typical for the diagrams of the sort of Figs. 5 c, d. For example, they occur in the calculation of the polarization vector when an electron is scattered by the Coulomb field in the second Born approximation. Terms proportional to $1/\Delta$ cancel in the spin independent part of the matrix elements (16) in the reducible and irreducible parts independently. However, in the spin dependent terms (17), (18) $1/\Delta$ cancels only in the sum of the irreducible and reducible parts in the nonrelativistic limit.

Taking into account all mentioned above, we write out the matrix elements for the diagrams 5 c, d. The reducible parts are

$$\mathcal{F}_N(\tilde{\gamma}) = -7 + 15 \tilde{\gamma}^2, \quad \mathcal{F}_{SO}(\tilde{\gamma}) = \frac{47}{16} \tilde{\gamma} + \frac{7}{8} \frac{\tilde{\gamma}}{\tilde{\gamma}^2 - 1}, \quad (30)$$

$$\mathcal{F}_{SS}^{(1)}(\tilde{\gamma}) = -4(1 - 2 \tilde{\gamma}^2) + \frac{31}{12} + \frac{1}{2} \frac{1}{\tilde{\gamma}^2 - 1}, \quad \mathcal{F}_{SS}^{(2)}(\tilde{\gamma}) = 4(1 - 2 \tilde{\gamma}^2) - \frac{77}{24} - \frac{1}{2} \frac{1}{\tilde{\gamma}^2 - 1}. \quad (31)$$

Since in the expressions (30), (31) we take the limit $t \rightarrow 0$, the singularity $1/\Delta$ turns to the singularity $1/(\tilde{\gamma}^2 - 1)$. Terms not proportional to $1/(\tilde{\gamma}^2 - 1)$ can be again represented in the form of an effective operator

$$\mathcal{L}_{ff}^{(cd)} = \sqrt{-g} \log |q^2| G^2 \left(\frac{15}{2} T^{\mu\nu} T_{\mu\nu} - \frac{7}{2} T^2 \right) + \frac{17}{2} \hat{O}_2 + \frac{31}{12} \hat{O}_3 - \frac{77}{24} \hat{O}_4. \quad (32)$$

The irreducible parts are

$$\mathcal{F}_N(\tilde{\gamma}) = -16 \tilde{\gamma} (2\tilde{\gamma}^2 - 1) \mathcal{W}_A(\tilde{\gamma}) + \frac{2}{3} (1 - 2\tilde{\gamma}^2)^2 \mathcal{W}_S(\tilde{\gamma}), \quad (33)$$

$$\mathcal{F}_{SO}(\tilde{\gamma}) = - \left(4\tilde{\gamma}^2 + \frac{1}{2} + \frac{7}{8} \frac{1}{\tilde{\gamma}^2 - 1} \right) \mathcal{W}_A(\tilde{\gamma}) + \frac{\tilde{\gamma}}{6} (2\tilde{\gamma}^2 - 1) \mathcal{W}_S(\tilde{\gamma}), \quad (34)$$

$$\mathcal{F}_{SS}^{(1)}(\tilde{\gamma}) = \left(2\tilde{\gamma} - 8\tilde{\gamma}^3 - \frac{1}{2} \frac{\tilde{\gamma}}{\tilde{\gamma}^2 - 1} \right) \mathcal{W}_A(\tilde{\gamma}) + \frac{1}{6} (1 - 2\tilde{\gamma}^2)^2 \mathcal{W}_S(\tilde{\gamma}), \quad (35)$$

$$\mathcal{F}_{SS}^{(2)}(\tilde{\gamma}) = \left(-2\tilde{\gamma} + 8\tilde{\gamma}^3 + \frac{1}{2} \frac{\tilde{\gamma}}{\tilde{\gamma}^2 - 1} \right) \mathcal{W}_A(\tilde{\gamma}) + \left(-\frac{1}{6} (2\tilde{\gamma}^2 - 1)^2 + \frac{1}{12} \right) \mathcal{W}_S(\tilde{\gamma}), \quad (36)$$

where functions $\mathcal{W}_S(\tilde{\gamma})$, $\mathcal{W}_A(\tilde{\gamma})$ are defined in Appendix by Eqs. (74), (75). As noted above, the terms singular in the $\tilde{\gamma} \rightarrow 1$ limit cancel in the sum of the reducible (30), (31) and irreducible (34)-(36) parts

$$\mathcal{F}_{SO}(1) = -\frac{13}{16}, \quad \mathcal{F}_{SS}^{(1)}(1) = \frac{7}{12}, \quad \mathcal{F}_{SS}^{(2)}(1) = -\frac{9}{8}. \quad (37)$$

4 Quantum corrections to spin effects

Summing up all the contributions we get the following scattering amplitudes. The terms not containing $S_\mu^{1,2}$ explicitly are

$$\mathcal{A}_N(q) = \frac{41}{5} G^2 m_1 m_2 \frac{(\bar{u}_1 u_1)}{2m_1} \frac{(\bar{u}_2 u_2)}{2m_2} \log |q^2|. \quad (38)$$

The terms proportional to the first power of $S_\mu^{1,2}$ are

$$\mathcal{A}_{SO}(q) = \frac{8}{5} i G^2 \epsilon_{\alpha\beta\gamma\delta} q^\alpha P_1^\beta P_2^\gamma \left(\frac{S_2^\delta}{2m_2^2} \frac{(\bar{u}_1 u_1)}{2m_1} + \frac{S_1^\delta}{2m_1^2} \frac{(\bar{u}_2 u_2)}{2m_2} \right) \log |q^2|. \quad (39)$$

At last, the part containing terms quadratic in spins is

$$\mathcal{A}_{SS} = \frac{G^2}{4m_1 m_2} \left(\frac{4}{15} (q \cdot S_1) (q \cdot S_2) + \frac{11}{40} t (S_1 \cdot S_2) \right) \log |q^2|. \quad (40)$$

Using the nonrelativistic expansions of the bispinor amplitudes of the scattering particles

$$\frac{1}{2m_{1,2}} \bar{u}(p_{1,2} \mp q) u(p_{1,2}) = 1 \pm \frac{i}{2m_{1,2}^2} \epsilon_{ijk} \mathbf{q}^i \mathbf{p}_{1,2}^j \mathbf{s}_{1,2}^k + \dots \quad (41)$$

$$\frac{1}{2m_{1,2}} S_{1,2}^\delta = \frac{1}{2m_{1,2}} \bar{u}(p_{1,2} \mp q) \gamma_5 \gamma^\delta \bar{u}(p_{1,2}) \approx (0, 2\mathbf{s}_{1,2}) \quad (42)$$

$$\frac{i}{2m_{1,2}^2} \epsilon_{\alpha\beta\gamma\delta} q^\alpha P_1^\beta P_2^\gamma S_{1,2}^\delta = \pm 8i \epsilon_{nkm} \mathbf{q}^n \mathbf{p}_{2,1}^k \mathbf{s}_{1,2}^m \mp 8i \frac{m_{2,1}}{m_{1,2}} \epsilon_{nkm} \mathbf{q}^n \mathbf{p}_{1,2}^k \mathbf{s}_{1,2}^m, \quad (43)$$

yields the amplitudes in the momentum space:

$$\mathcal{A}_N(\mathbf{q}) = \frac{41}{5} G^2 m_1 m_2 \log \mathbf{q}^2 \quad (44)$$

$$\mathcal{A}_{SO}(\mathbf{q}) = \frac{87}{10} G^2 \left(-i \frac{m_2}{m_1} \epsilon_{ijk} \mathbf{q}^i \mathbf{p}_1^j \mathbf{s}_1^k + i \frac{m_1}{m_2} \epsilon_{ijk} \mathbf{q}^i \mathbf{p}_2^j \mathbf{s}_2^k \right) \log \mathbf{q}^2, \quad (45)$$

$$+ \frac{64}{5} G^2 \left(i \epsilon_{nkm} \mathbf{q}^n \mathbf{p}_2^k \mathbf{s}_1^m - i \epsilon_{nkm} \mathbf{q}^n \mathbf{p}_1^k \mathbf{s}_2^m \right) \log \mathbf{q}^2, \quad (46)$$

$$\mathcal{A}_{SS}(\mathbf{q}) = G^2 \left(\frac{16}{15} (\mathbf{q} \cdot \mathbf{s}_1) (\mathbf{q} \cdot \mathbf{s}_2) + \frac{11}{10} \mathbf{q}^2 (\mathbf{s}_1 \cdot \mathbf{s}_2) \right) \log \mathbf{q}^2. \quad (47)$$

With the expressions (62) from Appendix we obtain the corrections in the coordinate space. First of all, according to [18], [20], the power quantum correction to the Newton law is the same as for scalar particles (1)

$$U_N = -\frac{41}{10} \frac{\hbar G^2 m_1 m_2}{c^3 \pi r^3}. \quad (48)$$

The quantum correction to the gravitational spin-orbit interaction is

$$U_{SO} = \frac{261}{20} \frac{\hbar G^2}{c^5 \pi r^5} \left(\frac{m_2}{m_1} \mathbf{s}_1 + \frac{m_1}{m_2} \mathbf{s}_2 \right) \cdot \mathbf{l}, \quad (49)$$

$$\mathbf{l}^i = \epsilon^{ijk} (\mathbf{r}_1 - \mathbf{r}_2)_j \mathbf{p}_k, \quad (50)$$

where $\mathbf{p} = \mathbf{p}_1 = -\mathbf{p}_2$ is the relative momentum.

The quantum correction to the Lense-Thirring effect is

$$U_{LT} = \frac{96}{5} \frac{\hbar G^2}{c^5 \pi r^5} (\mathbf{s}_1 + \mathbf{s}_2) \cdot \mathbf{l}. \quad (51)$$

The quantum correction to the interaction of the two internal angular momenta $\mathbf{s}_i$, i.e., to the gravitational spin-spin interaction is

$$U_{SS} = \frac{\hbar G^2}{c^5 \pi r^5} \left(8 (\mathbf{n} \cdot \mathbf{s}_1) (\mathbf{n} \cdot \mathbf{s}_2) + \frac{5}{2} (\mathbf{s}_1 \cdot \mathbf{s}_2) \right). \quad (52)$$

5 Discussion and conclusions

Hence, as expected, the quantum corrections to the classical Hamiltonian (49), (51), (52) for a spinor particle differ from the full sum of the similar corrections obtained in [20] for a rotating compound body with scalar constituents.

This difference is associated with the irreducible part as well as with the reducible one that can be represented in the form of an effective operator (see (2), (21), (25), (27), (32))

$$\mathcal{L} = \mathcal{L}_0 - 8\hat{O}_1 + 16\hat{O}_2 + \frac{31}{12} \hat{O}_3 - \frac{77}{24} \hat{O}_4, \quad (53)$$

where the new operators $\hat{O}_i$ take part.

Following are several points concerning the operator (53). As demonstrated in [20], after averaging over the spins, the Compton amplitude of fermion-graviton scattering and the scalar-graviton one coincide with the required accuracy. In particular, it means that the effects linear in spin (see expressions (49), (51)) for fermion-fermion and fermion-scalar scattering are the same. However, we see that the analysis is complicated by the singularity $1/\Delta$. Therefore, it is not unreasonable to make sure that the separation into the reducible and the irreducible parts is the same as in Eqs.(27), (30), (32), (33) for the case of fermion scattering by a scalar particle (evidently, there is only a contribution of one of the particles under the summation sign in the operators (32), (33) in this case). We use the effective vertices derived in [20] for scalar particles. The diagrams Figs. 5 a, b make the following contribution to the spin effects:

$$\mathcal{F}_N(\tilde{\gamma}) = 5 - 16\tilde{\gamma}^2, \quad \mathcal{F}_{SO}(\tilde{\gamma}) = -\frac{3}{4} \tilde{\gamma}. \quad (54)$$

The reducible part of the diagrams Figs. 5 c, d yields

$$\mathcal{F}_N(\tilde{\gamma}) = -9 + 16\tilde{\gamma}^2, \quad \mathcal{F}_{SO}(\tilde{\gamma}) = \frac{11}{4} \tilde{\gamma} + \frac{7}{8} \frac{\tilde{\gamma}}{\tilde{\gamma}^2 - 1}. \quad (55)$$

Note that the sum of (54) and (55) agree with the sum of corresponding expressions (26) and (30). The irreducible part is also equal to (34). Therefore the singularities $1/(\tilde{\gamma}^2 - 1)$ cancel in the same way as in the case of fermion-fermion scattering.

Thus, the effective operator (53) is universal for scattering of scalar particles and particles of spin 1/2.

Being interested in the quantum corrections to spin effects at the leading order in $1/c$ (where c is the speed of light), we considered only the matrix elements like (16)-(18). Formally, it is possible to consider one more matrix element at the same order in q/m_i

$$\mathcal{M}(s, t) = \mathcal{F}(s, t) t \frac{(P_1 S_2)}{m_2} \frac{(P_2 S_1)}{m_1}, \quad (56)$$

that however does not contribute to the spin effects at the leading order in $1/c$. Nevertheless, it is possible to make sure that the contribution of the diagrams Figs. 3, 4, 5 a, b and the reducible part of the diagram Figs. 5 c, d to the the form factor $\mathcal{F}(s, t)$ can also be described by the operator (53). This fact is trivial for the diagrams presented in Fig. 3 since their contribution originally contains the energy-momentum tensors of interacting particles. Contribution of the diagrams Fig. 4 is

$$\mathcal{F}(\tilde{\gamma}) = -\frac{\tilde{\gamma}}{4}, \quad (57)$$

that corresponds to the contribution of the operator (25). The diagrams Figs. 5 a, b yield

$$\mathcal{F}(\tilde{\gamma}) = 0, \quad (58)$$

that is also equal to the contribution of the corresponding operator (27). The reducible contribution of the diagrams Figs.(5)c, d is

$$\mathcal{F}_{\text{red}}(\tilde{\gamma}) = 2\tilde{\gamma} - \frac{33\tilde{\gamma}}{16(1-\tilde{\gamma}^2)} + \frac{\tilde{\gamma}}{4(1-\tilde{\gamma}^2)^2}. \quad (59)$$

Note that the terms not containing $1/(1-\tilde{\gamma}^2)$ also correspond to the contribution of the operator (32). Hence, the operator (53) describes all the contributions to the matrix element (56) correctly. Terms in the expression (59) singular in the $\tilde{\gamma} \rightarrow 1$ limit cancel with the irreducible part of the diagrams Figs. 5c, d in the same manner as for the matrix elements (17), (18):

$$\mathcal{F}_{\text{irred}}(\tilde{\gamma}) = \left(\frac{\tilde{\gamma}^3}{6} + \frac{\tilde{\gamma}}{48(1-\tilde{\gamma}^2)} \right) \mathcal{W}_S + \left(-2(1+\tilde{\gamma}^2) + \frac{9}{4(1-\tilde{\gamma}^2)} - \frac{1}{4(1-\tilde{\gamma}^2)^2} \right) \mathcal{W}_A, \quad (60)$$

$$\lim_{\tilde{\gamma} \rightarrow 1} (\mathcal{F}_{\text{red}}(\tilde{\gamma}) + \mathcal{F}_{\text{irred}}(\tilde{\gamma})) = -\frac{19}{40}. \quad (61)$$

I would like to thank I.B. Khriplovich and A.A. Pomeransky for their helpful comments and discussions. The investigation was supported by the Russian Foundation for Basic Research through Grant No. 05-02-16627-a and by Institute für Theoretische Physik-II Ruhr-Universität through Graduiertenkolleg "Physik der Elementarteilchen an Beschleunigern und im Universum".

Appendix

To calculate the correction to spin-dependent interactions, it is needed to consider the following Fourier transforms:

$$\begin{aligned} \int \log \mathbf{q}^2 e^{-i\mathbf{q}\cdot\mathbf{r}} \frac{d^3\mathbf{q}}{(2\pi)^3} &= -\frac{1}{2\pi r^3}, & \int (\mathbf{q}_i \mathbf{q}_j \log \mathbf{q}^2) e^{-i\mathbf{q}\cdot\mathbf{r}} \frac{d^3\mathbf{q}}{(2\pi)^3} &= \frac{3}{2\pi r^5} (5 \mathbf{n}_i \mathbf{n}_j - \delta_{ij}), \\ \int (-i\mathbf{q}_i \log \mathbf{q}^2) e^{-i\mathbf{q}\cdot\mathbf{r}} \frac{d^3\mathbf{q}}{(2\pi)^3} &= \frac{3}{2\pi} \frac{\mathbf{r}_i}{r^5}, & \int (\mathbf{q}^2 \log \mathbf{q}^2) e^{-i\mathbf{q}\cdot\mathbf{r}} \frac{d^3\mathbf{q}}{(2\pi)^3} &= \frac{3}{\pi r^5}. \end{aligned} \quad (62)$$

According to the choice $\langle 1(p_1 - q), 2(p_2 + q) | 1(p_1), 2(p_2) \rangle$ of kinematics and the sign in the exponent of the Fourier transforms, the vector $\mathbf{r}$ is directed from the second body to the first one, i.e., $\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2$.

Logarithmic parts of the integrals occurring are given below, using the notation $L = -\frac{i}{(4\pi)^2} \log |q^2|$:

$$I = \int \frac{d^D k}{(2\pi)^D} \frac{1}{k^2(k-q)^2} = L + \dots \quad (63)$$

$$I_\mu = \int \frac{d^D k}{(2\pi)^D} \frac{k_\mu}{k^2(k-q)^2} = \frac{q_\mu}{2} L + \dots \quad (64)$$

$$I_{\mu\nu} = \int \frac{d^D k}{(2\pi)^D} \frac{k_\mu k_\nu}{k^2(k-q)^2} = \frac{1}{3} \left(q_\mu q_\nu - \frac{q^2}{4} \delta_{\mu\nu} \right) L + \dots \quad (65)$$

$$J = \int \frac{d^D k}{(2\pi)^D} \frac{1}{k^2(k-q)^2((p-k)^2-m^2)} = \frac{L}{2m^2} + \dots \quad (66)$$

$$J_\mu = \int \frac{d^D k}{(2\pi)^D} \frac{k_\mu}{k^2(k-q)^2((p-k)^2-m^2)} = -\frac{p_\mu - q_\mu}{2m^2} L + \dots \quad (67)$$

$$J_{\mu\nu} = \int \frac{d^D k}{(2\pi)^D} \frac{k_\mu k_\nu}{k^2(k-q)^2((p-k)^2-m^2)} = \quad (68)$$

$$\frac{L}{4m^2} \left\{ \frac{q^2}{m^2} p_\mu p_\nu - (p_\mu q_\nu + p_\nu q_\mu) + 2q_\mu q_\nu - \frac{q^2}{2} g_{\mu\nu} \right\} + \dots \quad (69)$$

For the diagrams Fig. 5 c, d to be calculated, it is necessary to consider integrals with four Feynman propagators with the infrared divergency regularized by the "mass" of graviton λ . The imaginary part of the diagram Fig.5 c will be omitted because it does not contribute to the potential of the two-body interaction. It corresponds to the second Born approximation. The real part of the integral for the diagram Fig.5 c is

$$\begin{aligned} K^{(s)} &= \int \frac{d^D k}{(2\pi)^D} \frac{1}{(k^2 + i0) ((q-k)^2 + i0) ((p_1-k)^2 - m_1^2 + i0) ((p_2+k)^2 - m_2^2 + i0)} \\ &= -\frac{i}{(4\pi)^2} \frac{1}{m_1 m_2 t} \log \left| \frac{t}{\lambda^2} \right| \frac{2 m_1 m_2}{\sqrt{(s-m_+^2)(s-m_-^2)}} \log \frac{\sqrt{s-m_-^2} - \sqrt{s-m_+^2}}{\sqrt{s-m_-^2} + \sqrt{s-m_+^2}} + \dots \\ &= -\frac{i}{(4\pi)^2} \frac{1}{m_1 m_2 t} \log \left| \frac{t}{\lambda^2} \right| \mathcal{K}_s(s) + \dots, \end{aligned} \quad (70)$$

where notations $m_+ = m_1 + m_2$, $m_- = m_1 - m_2$ were introduced, and dependence on the variable s denoted as the function $\mathcal{K}_s(s)$. To simplify expressions it is convenient to introduce a new variable $\tilde{\gamma} = (s - m_1^2 - m_2^2)/(2m_1 m_2)$.

$$\mathcal{K}_s(s) = \frac{1}{\sqrt{\tilde{\gamma}^2 - 1}} \log \left(\tilde{\gamma} - \sqrt{\tilde{\gamma}^2 - 1} \right). \quad (71)$$

In the diagram Fig.5 d the same integral can be derived by analytic continuation of all the expressions from the s - to the u -channel ($t = q^2, u = 2m_1^2 + 2m_2^2 - s - t$):

$$\mathcal{K}_u(u) = -\mathcal{K}_s(2m_1^2 + 2m_2^2 - s - t) = \frac{2 m_1 m_2}{\sqrt{(m_+^2 - u)(m_-^2 - u)}} \log \frac{\sqrt{m_+^2 - u} + \sqrt{m_-^2 - u}}{\sqrt{m_+^2 - u} - \sqrt{m_-^2 - u}}. \quad (72)$$

In the limit when the momentum transfer q^μ goes to zero the function $\mathcal{K}_u(u)$ can be expressed as a function of $\mathcal{K}_s(s)$ up to terms linear in t :

$$\mathcal{K}_u(u) \approx -\mathcal{K}_s(s) + \frac{t}{2m_1m_2} \frac{\tilde{\gamma} \mathcal{K}_s(s) + 1}{\tilde{\gamma}^2 - 1}. \quad (73)$$

In the course of the calculation, it is convenient to express amplitudes as a half-sum and half-difference of the functions $\mathcal{K}_s(s)$ and $\mathcal{K}_u(u)$:

$$\mathcal{W}_S(\tilde{\gamma}) = -\lim_{t \rightarrow 0} \frac{6m_1m_2}{t} \frac{1}{2} (\mathcal{K}_s(s) + \mathcal{K}_u(u)) = -3 \frac{\tilde{\gamma} \mathcal{K}_s(s) + 1}{\tilde{\gamma}^2 - 1} \quad (74)$$

$$\mathcal{W}_A(\tilde{\gamma}) = \lim_{t \rightarrow 0} \frac{1}{2} (\mathcal{K}_u(u) - \mathcal{K}_s(s)) = -\mathcal{K}_s(s), \quad (75)$$

the functions $\mathcal{W}_S(\tilde{\gamma})$, $\mathcal{W}_A(\tilde{\gamma})$ are normalized to unity in the nonrelativistic limit, i.e., $\mathcal{W}_S(1) = \mathcal{W}_A(1) = 1$.

Vector and tensor integrals with four propagators are considered with respect to the relations already obtained (63)-(70), (72). As an example, consider an integral ($P_1 = 2p_1 - q$, $P_2 = 2p_2 + q$):

$$K_\mu^{(s)} = \int \frac{d^D k}{(2\pi)^D} \frac{k_\mu}{(k^2 - \lambda^2)((k - q)^2 - \lambda^2)(k^2 - 2p_1 k)(k^2 + 2p_2 k)} =$$

$$= AP_{1\mu} + BP_{2\mu} + Cq_\mu. \quad (76)$$

$$(77)$$

Using the following relations for the logarithmic parts, we obtain the coefficients A , B , C :

$$q^\mu K_\mu^{(s)} = \frac{q^2}{2} K^{(s)} \quad (78)$$

$$P_1^\mu K_\mu^{(s)} = -J_2 - \frac{q^2}{2} K^{(s)} \quad (79)$$

$$P_2^\mu K_\mu^{(s)} = J_1 + \frac{q^2}{2} K^{(s)}, \quad (80)$$

here J_1 and J_2 are the integrals (66) with the propagators of the first and the second particles, respectively;

$$A = \frac{4s}{\Delta} \left[\left(J_1 + \frac{q^2}{2} K^{(s)} \right) (P_1 P_2) + \left(J_2 + \frac{q^2}{2} K^{(s)} \right) P_2^2 \right] \quad (81)$$

$$B = -\frac{4s}{\Delta} \left[\left(J_2 + \frac{q^2}{2} K^{(s)} \right) (P_1 P_2) + \left(J_1 + \frac{q^2}{2} K^{(s)} \right) P_1^2 \right] \quad (82)$$

$$C = \frac{1}{2} K^{(s)}, \quad (83)$$

where we introduce the following notation

$$\Delta = \frac{1}{4s} \left((P_1 P_2)^2 - P_1^2 P_2^2 \right) \quad s = (p_1 + p_2)^2. \quad (84)$$

Tensor integrals can be considered in the same way.

References

- [1] A.F. Radkowski, Ann. of Phys. **56**, 319 (1970).
- [2] D.M. Capper, M.J. Duff, L. Halpern, Phys. Rev. **D10**, 461 (1974).
- [3] D.M. Capper, M.J. Duff, Nucl. Phys. **B44**, 146 (1974).
- [4] M.J. Duff, J.T. Liu, Phys.Rev.Lett. **85** (2000) 2052;
hep-th/0003237.
- [5] D.M. Capper, G. Leibbrandt, M. Ramon Medrano, Phys. Rev. **D8**, 4320 (1973).
- [6] M.J. Duff, Phys. Rev. **D9**, 1837 (1973).
- [7] G. 't Hooft, M. Veltman, Ann. Inst. H. Poincare **A20**, 69 (1974).
- [8] D.G. Boulware, S. Deser, Ann. Phys. **89**, 193 (1975).
- [9] J.F. Donoghue, Phys. Rev. Lett. **72**, 2996 (1994); gr-qc/9310024.
- [10] J.F. Donoghue, Phys. Rev. **D50**, (1994) 3874; gr-qc/9405057.
- [11] J.F. Donoghue, Introduction to the effective field theory description of gravity,
in *Advanced School on Effective Theories*, ed. by F. Cornet and M.J. Herrero
(World Scientific, Singapore, 1996); gr-qc/9512024.
- [12] J.F. Donoghue, Perturbative dynamics of quantum general relativity,
in *Proceedings of the Eighth Marcel Grossmann Meeting on General Relativity*,
ed. by Tsvi Piran and Remo Ruffini (World Scientific, Singapore, 1999);
gr-qc/9712070.
- [13] I.J. Muzinich, S. Vokos, Phys. Rev. **D52**, 3472 (1995); hep-th/9501083.
- [14] A. Akhundov, S. Belucci, A. Shiekh, Phys. Lett. **B395**, 19 (1998); gr-qc/9611018.
- [15] H. Hamber, S. Liu, Phys. Lett. **B357**, 51 (1995); hep-th/9505182.
- [16] D. Dalvit, F.D. Mazzitelli, Phys. Rev. **D56**, 7779 (1997); hep-th/9708102.
- [17] I.B. Khriplovich, G.G. Kirilin, Zh. Eksp. Teor. Fiz. **109**, 1139 (2002);
[Sov. Phys. JETP **95**, 981 (2002)]; gr-qc/0207118.
- [18] N.E.J. Bjerrum-Bohr, J.F. Donoghue, B.R. Holstein, Phys. Rev. **D67**, (2003) 084033;
gr-qc/0211072.
- [19] N.E.J. Bjerrum-Bohr, J.F. Donoghue, B.R. Holstein, Phys. Rev. **D68**, (2003) 084003;
gr-qc/0211071.
- [20] I.B. Khriplovich, G.G. Kirilin, Zh. Eksp. Teor. Fiz. **125**, 1219 (2004);
[Sov. Phys. JETP **98**, 1063 (2004)]; gr-qc/0207118.
- [21] I.B. Khriplovich, A.A. Pomeransky, Zh. Eksp. Teor. Fiz. **113**, 1537 (1998);
[Sov. Phys. JETP **86**, 839 (1998)]; gr-qc/9710098.